arXiv:1112.0531v1 [math.DS] 2 Dec 2011

A separation theorem for entire transcendental maps

Anna Miriam Benini [*]
CRM Ennio de Giorgi
Piazza dei Cavalieri 3
56100 Pisa, Italy
ambenini@gmail.com

Núria Fagella[†]
Dept. de Mat. Aplicada i Anàlisi
Univ. de Barcelona, Gran Via 585
08007 Barcelona, Spain
fagella@maia.ub.es

December 5, 2011

Abstract

We study the distribution of periodic points for a certain class of holomorphic functions. An interior periodic point of period p is a periodic point which is not the landing point of any periodic ray of period less or equal than p, like for example an attracting cycle or a Cremer cycle. In polynomial dynamics, it is known by a theorem of Goldberg and Milnor that periodic rays, together with their landing points, separate the plane into regions each containing exactly one interior periodic point or a parabolic immedite basin. We show an analogue of this theorem for a wide class of transcendental entire functions (those with a bounded set of singular values and finite order, or composition thereof), under the assumption that periodic rays land. This result has many corollaries. Among them, it follows that there cannot be Cremer points on the boundary of Siegel disks; that in the absence of wandering domains "hidden components" of a Siegel disk have to be preperiodic to the Siegel disk itself; or that there cannot be infinitely many attracting cycles of period less than any finite constant.

1 Introduction

Given a holomorphic map $f : \mathbb{C} \to \mathbb{C}$, we are interested in the dynamical system generated by the iterates of f. In this setup, there is a dynamically meaningful partition of the phase space, into two completely invariant subsets: the *Fatou set*, where the dynamics are stable, and the *Julia set* where they are chaotic. More precisely, the Fatou set is defined as the open set

$$\mathcal{F}(f) := \{z \in \mathbb{C}; \{f^n\} \text{ is normal in a neighborhood of } z\},$$

and the *Julia set* $J(f)$ as is its complement. Another dynamically interesting set is the set of escaping points or *escaping set*

$$I(f) := \{z \in \mathbb{C};\ f^n(z) \to \infty, \text{ as } n \to \infty\}.$$

---

[*] Partially supported by the EU network MRTN-CT-2006-035651 CODY

[†] Partially supported by the catalan grant 2009SGR-792, by the spanish grants MTM-2008-01486 and MTM2006- 05849 Consolider (including a FEDER contribution) and by the european network MRTN-CT-2006-035651-2-CODY.

The relation between them is that, in general, $J(f) = \partial I(f)$, although for some classes of functions $I(f) \subset J(f)$ ([EL]) and hence $J(f) = \overline{I(f)}$.

In this paper we are mainly concerned with *entire transcendental* maps (abbreviated transcendental maps), i.e., those entire maps for which infinity is an essential singularity. There are several essential differences between the dynamics of transcendental maps and that of polynomials, coming from the very different behavior of iterates in a neighborhood of infinity. For example, while the Julia set of polynomials is always a compact set disjoint from $I(f)$, its analogous for transcendental maps is always unbounded, an contains infinitely many continua in $I(f)$. In fact, the escaping set is part of the Fatou set in the case of polynomials, while it is part of the Julia set for transcendental maps.

Another relevant difference concerns the singularities of the inverse function, which always play a crucial role in holomorphic dynamics. For polynomials (or rational maps) all branches of f^{-1} are well defined in a neighborhood of any point, with the exception of *critical values*, that is, $v = f(c)$ where c is a zero of f' or *critical point.* If f is entire, one more type of singularity is allowed, namely the *asymptotic values*, or points $a \in \mathbb{C}$ for which there is a curve $\gamma(t)$ such that $|\gamma(t)| \to \infty$ as $t \to \infty$, and $f(\gamma(t)) \to a$ as $t \to \infty$. In some loose sense, asymptotic values have some of their preimages at infinity.

A holomorphic function is a covering outside the set $S(f)$ of *singular values* of f, where

$$\mathcal{S}(f) := \overline{\{z \in \mathbb{C};\ z \text{ is an asymptotic or critical value for } f\}}.$$

Observe that rational maps, and in particular polynomials, have a finite number of singular values while transcendental maps may have infinitely many such singularities. This is in fact part of the reason why for polynomials, every component of the Fatou set (also called *Fatou component*) is preperiodic to a periodic attracting or parabolic basin or to a periodic rotation domain (Siegel disk). Instead, transcendental maps allow for additional types of Fatou components like *wandering components* (i.e., not preperiodic) or *Baker domains* (i.e., components for which all points tend to infinity under iteration). However, there exist certain natural classes of transcendental maps for which the dynamics is in some sense similar to that of polynomials. One such class is the Eremenko-Lyubich class

$$\mathcal{B} = \{f : \mathbb{C} \to \mathbb{C} \text{ entire transcendental; } S(f) \text{ is bounded}\}.$$

For functions in class $\mathcal{B}$, the escaping set is a subset of the Julia set (see [EL], Theorem 1) and therefore there are neither Baker domains nor wandering domains in which the iterates tend to infinity. It is not known actually whether they may have other types of wandering domains. If $S(f)$ is finite, then f is said to be of *finite type* and has no wandering domains of any kind ([EL], [GK]).

Recall that a function is of *finite order* if $\log\log|f(z)|$ is of the order of $\log|z|$ as $|z| \to \infty$. For example, $f(z) = e^{z^p}$ has order p. In this paper we will restrict ourselves to the class

$$\widehat{\mathcal{B}} = \{f = g_1 \circ \ldots \circ g_k;\ k \geq 1,\ g_i \in \mathcal{B} \text{ and } g_i \text{ has finite order}\}.$$

Observe that class $\widehat{\mathcal{B}}$ is closed under composition, as is $\mathcal{B}$, but not the class of functions of finite order.

The reason why we restrict ourselves to class $\widehat{\mathcal{B}}$, is because it is the most general class for which *(dynamic) rays* or *hairs* have been proven to exist. Informally, a ray is a maximal unbounded injective curve $g : (0, \infty) \to I(f)$ such that all its iterates are also unbounded curves with the same properties (see Section 2 for a precise definition). We say that a ray *lands* at $z_0 \in \mathbb{C}$ if $g(t) \to z_0$ as $t \to 0$. By construction, rays are always pairwise disjoint.

For polynomials, rays foliate the attracting basin of infinity and belong to the Fatou set (see for example [DH], [Mi]). They are in one to one correspondence with straight rays in the complement of $\overline{\mathbb{D}}$ via Böttcher coordinates, and therefore its dynamics is governed by $\theta \mapsto d\theta$, where θ is the angle that identifies the ray and d is the degree of the polynomial (recall that a polynomial behaves like $z \mapsto z^d$ in a neighborhood of infinity).

For transcendental maps, when rays exist, they belong to the Julia set. The existence of rays, and their organization with respect to some symbolic dynamics was settled initially in [DT] for functions of finite type (with some extra technical condition). Recently in [R3S], these results have been extended to the class $\widehat{\mathcal{B}}$ (in fact, to a more general but less natural class than $\widehat{\mathcal{B}}$). We refer to Section 2 for background and precise statements. Of special importance for us will be the $p-$*periodic* rays, i.e., rays that come back to themselves after p iterates. As for periodic points, p is called the *period* of the ray, which we will in general not assume to be minimal. When the period is 1, the ray is called *invariant* or *fixed*.

Our goal in the present work is to prove some results concerning the relation between periodic rays, their landing points and the distribution of some special periodic points in the complex plane. More precisely, a $p-$periodic point is called an *interior* $p-$*periodic point* if there are no $p-$periodic rays landing at it.

Periodic points which are neither repelling nor parabolic (hence (super)attracting, Siegel or Cremer) are necessarily interior periodic points. However, repelling or parabolic $p-$periodic points can be interior periodic points as well, for example if they are landing points of rays of higher period or if they are not landing points of any ray.

To state our results we need to partially recall how the dynamics around a parabolic point is. If z_0 is a parabolic $p-$periodic point, an *attracting petal* at z_0 is defined an any open simply connected set U containing z_0 on its boundary, such that $f^p(\overline{U}) \subset U \cup \{z_0\}$. A *repelling petal* is an attracting petal for the branch of f^{-p} which fixes z_0. It is well known (see the Fatou Flower Theorem 6.2) that the number of attracting petals at z_0 equals its multiplicity as a fixed point of f, minus one. The number of repelling petals is the same and in fact, they distribute themselves alternatingly around z_0 forming a neighborhood of the parabolic point.

We will first restrict the attention to fixed points and fixed rays. We assume that all fixed rays land in $\mathbb{C}$, and in fact this will be a standing assumption for the whole paper. If two periodic rays land together at a common point, the curve they form together with their common landing point is called a *ray pair*. In contrast, we say that a ray *lands alone* if its landing point is not the landing point of any other ray.

Let Γ be the graph formed by fixed rays together with their endpoints. A connected component of $\mathbb{C} \setminus \Gamma$ is called a *basic region* for f.

The next theorem is proven by Goldberg and Milnor in [GM, Thm. 3.3].

Theorem 1.1 (GM Theorem). *Let f be a polynomial with connected Julia set. Then each* *basic region contains exactly one interior fixed point, or an invariant attracting parabolic petal.*

In the proof of this theorem, the finiteness of the degree of f is used in an essential way. In a suitable restriction of each basic region, a weakly polynomial-like map is constructed and the Lefschetz fixed point theory is used to count the number of critical points and fixed points.

The strategy as it is cannot be adapted to the transcendental setting, where the degree is hardly well defined on any restriction, and where the role of critical values can be carried out also by asymptotic values. However, we use Goldberg and Milnor's idea of doing a counting in the basic regions, and we manage to do it successfully by counting directly fixed points instead of going through critical points. In this way we prove a transcendental analog of the GM Theorem.

Theorem A (Separation theorem). Let $f \in \widehat{\mathcal{B}}$ such that all fixed rays land. Then there are finitely many basic regions, and each basic region contains exactly one interior fixed point, or an invariant attracting parabolic petal.

We remark that in the polynomial case, the hypothesis on connectivity of the Julia set was enough to ensure that all fixed rays landed. For transcendental functions, there is no necessary and sufficient condition to show that fixed rays land. There are however many cases in which this can be shown, for example if the postsingular set

$$\mathcal{P}(f) : \overline{\bigcup_{s \in \mathcal{S}(f), n \in \mathbb{N}} \{f^n(s)\}}$$

does not intersect the set of fixed rays (see [Fa], [Rel]).

Let us observe some facts which are obvious for polynomials but not at all obvious for transcendental maps. A polynomial of degree d always has $d-1$ fixed rays (fixed points of $\theta \mapsto d\theta$) and d fixed points. This means that there is always at least one interior fixed point, even if all rays lands alone (this would be the case with only one basic region). If certain rays land together, then more fixed points become interior fixed points as they cannot be landing points of any fixed ray. This basic counting, and the fact that there are only finitely many basic regions, is possible only because of the finiteness of the degree. In the transcendental case there are infinitely many fixed points and infinitely many rays, so a priori one might have infinitely many ray pairs, basic regions and interior fixed points. Theorem A already states that f has a finite number of those, but moreover, a similar global counting as in the polynomial case can be done.

Theorem B (Global counting). Let $f \in \widehat{\mathcal{B}}$ such that all fixed rays land. Let $G = \{g_1, \ldots, g_N\}$ be a collection of fixed rays satisfying that any ray $g \notin G$ lands alone. Then, there are exactly $N+1$ fixed points which are either landing points of some $g \in G$ or interior fixed points.

Like in the polynomial case, the statement about periodic points follows from the fixed case and from the fact that the class $\widehat{\mathcal{B}}$ is invariant under composition. Since this is the most general result and the one that is used in the applications, we state it as a theorem although it follows immediately from Theorem A.

Theorem C (Separation theorem for periodic rays). Let $f \in \widehat{\mathcal{B}}$ such that p-periodic rays of period land. Then there are finitely many basic regions for f^p, and any two interior p-periodic points can be separated by a periodic ray pair of the same period p.

The true power of the GM Theorem, and therefore of our separation theorems, resides in their remarkable number of corollaries. Some of them were not yet known for transcendental functions of any class. We summarize them in the following statement.

Corollary D. If f is a function in $\widehat{\mathcal{B}}$ whose periodic rays land,

1. There are only finitely many interior periodic points of any given period.
2. Any two periodic Fatou components can be separated by a periodic ray pair.
3. There are no Cremer points on the boundary of periodic Fatou components.
4. If z_0 is a parabolic point, any two of its attracting petals are separated by a ray pair of the same period. In particular, for each repelling petal there is at least one ray which lands at z_0.
5. For any given period p, there are only finitely many (possibly none) periodic points of period p which are landing points of more than one periodic ray. None of them is landing point of infinitely many rays of the same period.

Note that it follows from statement 1 that f cannot have infinitely many attracting cycles of period less or equal than any given constant, regardless of the presence of infinitely many singular values.

From the fact that Fatou components can be separated by periodic ray pairs we obtain the following additional corollary. Given an invariant Siegel disk Δ, we say that U is a hidden component of Δ if U is a bounded connected component of $\mathbb{C} \setminus \overline{\Delta}$.

Corollary E (Hidden components of Siegel disks). If $f \in \widehat{\mathcal{B}}$ has no wandering domains, any hidden component of a bounded invariant Siegel disk is preperiodic to the Siegel disk itself.

This article is structured as follows. In Section 2 we state and prove a result concerning the existence of fixed rays for functions in $\widehat{\mathcal{B}}$ and some expansion properties (Lemma 2.2). We call it the Structural Lemma and it is a consequence of results in [R3S]. Hence, we must introduce logarithmic coordinates and the setup of [R3S] to prove it. We also setup some additional symbolic dynamics to be able to transfer their statements to our dynamical plane. This setup has interest on its own and we hope that it can be used for other purposes as well. With the Structural Lemma in hand, in Section 3 we present some results about the topology and general distribution of fixed points in the plane. In Section 4 we introduce the argument principle and prove some homotopy lemmas used in the two following sections, devoted to the proof of Theorems B and A respectively. Finally, in Section 7 we prove corollaries D and E.

Acknowledgments

We are especially thankful to Arnaud Chéritat and to Pascale Roesch for sharing with us the preprint of their recent work on Siegel disks for polynomial with two critical values ([CR]),

from which comes the original idea that ended up generating this present paper. We thank Xavier Buff for helpful discussions and we are also thankful to the Complex Dynamics group in Universitat de Barcelona, and to the CODY Marie Curie Network thanks to which this project could be carried out.

2 Tools and preliminary results

In this section, we will first present the main features of the dynamical plane for $f \in \widehat{\mathcal{B}}$ and state a Structural Lemma (Lemma 2.2) which is needed for the rest of the paper. We will then prove it using results of [R3S] together with some study of the combinatorics of f. This last part of the section is necessary for the proof and interesting in itself, but will not be used later during the paper.

2.1 Structure of the dynamical plane and main technical lemma

Let $f \in \mathcal{B}$, and D be a Jordan domain containing $S(f), 0$ and $f(0)$. For simplicity we assume that the boundary of D is real analytic. Then the preimage $\mathcal{T}$ of $\mathbb{C} \setminus \overline{D}$ consists of countably many unbounded connected components $\{T_\alpha, \alpha \in \mathcal{A}\}$ called *tracts* of f, and $f : T_\alpha \to \mathbb{C} \setminus \overline{D}$ is a universal covering of infinite degree.

Observe that only finitely many tracts intersect any compact set because fibers are discrete. Therefore, we can choose a curve $\delta \subset \mathbb{C} \setminus (\overline{D} \cup \overline{\mathcal{T}})$ connecting $\overline{D}$ to infinity. The preimages $\{f^{-1}\delta\}$ subdivide each tract T_α into *fundamental domains* $F_{\alpha,i}$ with $i \in \mathbb{Z}$ on which $f : F_{\alpha,i} \to \mathbb{C} \setminus (\delta \cup \overline{D})$ is univalent (see Figure 1).



Figure 1: The disk D, the tracts of f and the fundamental domains.

When it is not relevant which tract the fundamental domains are contained in, or their position inside the tract, we call a fundamental domain simply F. Observe that for each F, because $f|_F$ is univalent, there is a well defined bijective inverse branch $f_F^{-1} : \mathbb{C}\setminus(\delta\cup\overline{D}) \to F$.

Definition 2.1 (Dynamic ray). A *(dynamic) ray* for f is an injective curve $g : (0,\infty) \to I(f)$ such that:

- $\lim\limits_{t\to\infty} |f^n(g(t))| = \infty \;\; \forall n \geq 0$;
- $\lim\limits_{n\to\infty} |f^n(g(t))| = \infty$ uniformly in $[t,\infty)$;
- $g(0,\infty)$ is maximal with these properties.

We say that a ray g is *asymptotically contained* in a tract F if $g(t) \in F$ for all t sufficiently large. Recall that a dynamic ray is g *periodic* (or m-periodic) if $f^m(g) \subset g$ for some $m \geq 1$, and *fixed* if $m = 1$. We say that a dynamic ray *lands* at a point $z_0 \in \mathbb{C}$ if $\lim g(t) = z_0$ as $t \to 0$.

The following lemma summarizes all of the results from this section that we shall use in the rest of the paper.

Lemma 2.2 (Structural Lemma). *Let $f \in \widehat{\mathcal{B}}$. Then*

(a) Tracts and fundamental domains have a cyclic order at infinity and can be labeled according to that order.

(b) For each fundamental domain F, there is a unique fixed dynamic ray g_F which is asymptotically contained in F. Conversely, any fixed ray is asymptotically contained in some fundamental domain.

(c) Let $\mathcal{F}$ be a finite collection of fundamental domains. Then for any R sufficiently large, the preimage of the circle C_R of radius R restricted to $\mathcal{F}$ is contained in the bounded connected component of $\mathbb{C} \setminus C_R$.

Each of the three remaining subsection is dedicated to the proof of each of the statements of Lemma 2.2.

2.2 Logarithmic coordinates and lift of f

For an entire transcendental function in class $\mathcal{B}$ we can consider logarithmic coordinates following [EL] (see also [R3S]). The logarithm is a well defined multivalued function on $\mathbb{C}\setminus\overline{D}$, so we can define the set $\widetilde{H} := \exp^{-1}(\mathbb{C}\setminus\overline{D})$, which contains a right half plane. Also $\exp^{-1}(\delta)$ is a countable family of curves $\widetilde{\delta}_i$ dividing $\widetilde{H}$ into semi-strips $\widetilde{S}_i$; without loss of generality we can label one of them $\widetilde{S}_0$ and find a branch L_0 of the logarithm mapping $\mathbb{C}\setminus(\overline{D}\cup\delta)$ into $\widetilde{S}_0$. If we define branches L_n of the logarithm as $L_n(w) = L_0(w) + 2\pi i n$, we obtain a consecutive labeling of the strips $\widetilde{S}_n$ as images of L_n. Call $\widetilde{\delta}_n, \widetilde{\delta}_{n+1}$ the preimages of δ bounding the semi-strip $\widetilde{S}_n$. Observe that the tracts do not contain the origin, so the branches of the logarithm are defined on the entire tracts (see Figure 2). The set $\widetilde{\mathcal{T}} := \exp^{-1}(\mathcal{T})$ then consists of infinitely many connected components $\widetilde{T}_{\alpha,i}$ which do not accumulate in any compact set; the set $\widetilde{T}_{\alpha,i}$ is defined as $L_i(T_\alpha)$ so an unbounded part of $\widetilde{T}_{\alpha,i}$ is always contained in $\widetilde{S}_i$.

The following properties hold (see [R3S]):



Figure 2: Logarithmic coordinates. The lift $\widetilde{f}$ is only defined on the tracts. Each of them is mapped conformally onto the set $\widetilde{H}$.

- $\widetilde{H}$ is a $2\pi i$ periodic domain containing a right half plane;
- every $\widetilde{T}_{\alpha,i}$ is an unbounded Jordan domain with real parts bounded from below, but unbounded from above (i.e., unbounded to the right);
- the $\widetilde{T}_{\alpha,i}$ accumulate only at infinity;
- $\widetilde{\mathcal{T}}$ is invariant under translation by $2\pi i$.

Lift of f

Because $f \circ \exp$ is a universal covering on each $\widetilde{T}_{\alpha,i}$, it is possible to lift f to a continuous function $\widetilde{f}:\widetilde{\mathcal{T}}\to\widetilde{H}$ which makes the following diagram commute.

$$\begin{tikzcd}
\widetilde{\mathcal{T}} \arrow[r, "\widetilde{f}"] \arrow[d, "\exp"'] & \widetilde{H} \arrow[d, "\exp"] \\
\mathcal{T} \arrow[r, "f"] & \mathbb{C}\setminus\overline{D}
\end{tikzcd}$$

On each of the infinitely many connected components $\widetilde{T}_{\alpha,i}$ of $\widetilde{\mathcal{T}}$, the lift $\widetilde{f}|_{\widetilde{T}_{\alpha,i}}$ is defined up to translation by multiples of $2\pi i$; so $\widetilde{f}:\widetilde{\mathcal{T}}\to\widetilde{H}$ is defined up to infinitely many constants. Let us fix any choice of $\widetilde{f}$ on each of the tracts $\widetilde{T}_{\alpha,0}$, and then choose the remaining constants so that $\widetilde{f}$ is $2\pi i$ periodic.

The preimages of $\{\widetilde{\delta}_j\}$ under $\widetilde{f}$ divide each $\widetilde{T}_{\alpha,i}$ into *fundamental domains* $\widetilde{F}_{\alpha,i,j}$ on which $\widetilde{f}$ is univalent. Choose the labeling of fundamental domains so that $\widetilde{f}(\widetilde{F}_{\alpha,i,j}))=\widetilde{S}_j$ for each α, i. Because we chose the lift $\widetilde{f}$ to be periodic, the labeling on the fundamental domains for $\widetilde{f}$ induces a well defined labeling for the fundamental domains for f, in such a way that $\exp(\widetilde{F}_{\alpha,i,j})=F_{\alpha,j}$ for each i (see Figure 3).


$\widetilde{\delta}_{n-1}$
$\widetilde{F}_{\alpha,n,2}$
$\widetilde{F}_{\alpha,n,1}$
$\widetilde{T}_{\alpha,n}$
$\widetilde{F}_{\alpha,n,0}$
T_α
exp
$F_{\alpha,2}$
$F_{\alpha,1}$
$\widetilde{F}_{\alpha,n,-1}$
$\widetilde{F}_{\alpha,n,-2}$
$F_{\alpha,0}$
$F_{\alpha,-1}$
$F_{\alpha,-2}$
$\widetilde{\delta}_n$
D
δ

Figure 3: Labeling of the fundamental domains.

Vertical order

Because the sets $\widetilde{T}_{\alpha,i}$ are unbounded to the right, each $\widetilde{T}_{\alpha,i}$ divides any right half plane into at least two connected components, one of which contains points with arbitrarily large imaginary part (that is said to be *above* $\widetilde{T}_{\alpha,i}$) and one which contains points with arbitrarily small imaginary part (that is said to be *below* $\widetilde{T}_{\alpha,i}$). This introduces a natural order $\prec$ on the $\widetilde{T}_{\alpha,i}$, called *vertical order at infinity.* We can now assume that $\widetilde{T}_{\beta,i} \prec \widetilde{T}_{\alpha,i}$ if and only if $\beta < \alpha$ (for a fixed i). Observe also that $\widetilde{T}_{\alpha,i} \prec \widetilde{T}_{\beta,j}$ whenever $i < j$; in this case, the order is just induced by branches of the logarithm. The vertical order on the $\widetilde{T}_{\alpha,i}$ induces via the exponential map a *cyclic order at infinity*(still denote by $\prec$) on the tracts T_α.

Observe that for any given tract T_α, the fundamental domains $F_{\alpha,i}$ contained in T_α also have a well defined cyclic order; this gives a total ordering on the set of fundamental domains.

This proves part (a) of the Structural Lemma 2.2.

2.3 Dynamic rays and their properties

Here we present a result in [R3S] about existence of dynamic rays for $\widetilde{f}$. Then, we prove a theorem about existence of dynamic rays for f corresponding to part (b) of the Structural Lemma 2.2.

We define the escaping set

$$I(\widetilde{f}) := \{z, \operatorname{Re} \widetilde{f}^n(z) \to \infty \text{ as } n \to \infty\}.$$

The following definition of ray tail and dynamic ray is taken from Definition 2.2 in [R3S].

Definition 2.3. Let $\widetilde{f}$ be the lift of a function f in class $\widehat{\mathcal{B}}$. A *ray tail* for $\widetilde{f}$ is an injective curve $\widetilde{g} : [t_0, \infty) \to I(\widetilde{f})$ such that $\lim_{t\to\infty} \operatorname{Re} \widetilde{f}^n(\widetilde{g}(t)) = +\infty$ for any $n > 0$ and such that $\operatorname{Re} \widetilde{f}^n(\widetilde{g}(t)) \to +\infty$ uniformly in t as $n \to \infty$.

Likewise ray tails can be defined for f. Observe that with this definition a dynamic ray for f becomes a maximal injective curve $g(t) : (0, \infty) \to I(f)$ such that $g|_{[t,\infty)}$ is a ray tail for every $t > 0$ large enough.

Let us define

$$J(\widetilde{f}) := \{\widetilde{z}, \widetilde{f}^n(\widetilde{z}) \text{ is defined for all } n\}.$$

Observe that $J(\widetilde{f})$ includes also fixed points, not only points escaping to infinity.

For points $\widetilde{z} \in J(\widetilde{f})$, we can naturally introduce symbolic dynamics using the labeling for the tracts (see [DT], [R3S],[Ba]): if $\widetilde{f}^n(\widetilde{z}) \in \widetilde{T}_{\alpha_n,i_n}$, we will say that $\widetilde{z}$ has *address*

$$\widetilde{s} := \begin{pmatrix} \alpha_0 \\ i_0 \end{pmatrix} \begin{pmatrix} \alpha_1 \\ i_1 \end{pmatrix} \ldots$$

We can then define the set

$$J_{\widetilde{s}} := \{\widetilde{z} \in J(\widetilde{f}), \widetilde{z} \text{ has address } \widetilde{s}\}$$

and introduce the left-sided shift map σ

$$\sigma : \begin{pmatrix} \alpha_0 \\ i_0 \end{pmatrix} \begin{pmatrix} \alpha_1 \\ i_1 \end{pmatrix} \ldots \mapsto \begin{pmatrix} \alpha_1 \\ i_1 \end{pmatrix} \begin{pmatrix} \alpha_2 \\ i_2 \end{pmatrix} \ldots .$$

From the definition of $J_{\widetilde{s}}$, it follows that $\widetilde{f}(J_{\widetilde{s}}) \subset J_{\sigma\widetilde{s}}$; in particular, if an address $\widetilde{s}$ is periodic of period m, we have that $\widetilde{f}^m(J_{\widetilde{s}}) \subset J_{\widetilde{s}}$.

The next theorem follows from results in [R3S].

Theorem 2.4 (**Transcendental rays**)**.** *Let* $\widetilde{f}$ *be the lift of a function* $f \in \widehat{\mathcal{B}}$*. Then for any address* $\widetilde{s}$*, if* $J_{\widetilde{s}} \cap I(\widetilde{f}) \neq \emptyset$ *it contains a unique unbounded arc consisting of escaping points, called the* ray tail *of address* $\widetilde{s}$ *for* $\widetilde{f}$*.*

Theorem 2.4 is not stated in this terms, so we include a sketch of proof using results and terminology from [R3S].

Proof. From Theorem 5.6 and Lemma 5.7 in [R3S], $\widetilde{f}$ satisfies a uniform linear head start condition (uniform head start and uniform bounded wiggling are equivalent by Proposition 5.4). From Proposition 4.5 (b), if the set

$$J_{\widetilde{s}}^{K} := \{z \in J_{\widetilde{s}}, \operatorname{Re} \widetilde{f}^{n}(z) \geq K \,\forall n \in \mathbb{N}\}$$

is non empty, then $J_{\widetilde{s}}$ has a unique unbounded component which is a closed arc to infinity. From Corollary 4.4, the mentioned arc consists of escaping points. Also, if $J_{\widetilde{s}}$ contains at least an escaping point, $J_{\widetilde{s}}^{K} \neq \emptyset$. □

Remark 2.5. If the address $\widetilde{s}$ contains only finitely many different symbols (for example, if it is periodic) standard arguments can be used to show that $J_{\widetilde{s}} \cap I(\widetilde{f}) \neq \emptyset$.

Now we show that each fundamental domain F asymptotically contains exactly one fixed ray. To do this we introduce addresses in the f-plane and a correspondence between addresses for f and addresses for $\widetilde{f}$. The content of this subsection is a natural consequence of the setup in [R3S], although it is not contained there.

Proposition 2.6 (**Fixed rays in fundamental domains**)**.** *For each fundamental domain* F*, there is a unique invariant dynamic ray* g_F *such that* $g_F(t) \subset F$ *for all sufficiently large* t*.*

To prove Proposition 2.6, we need to understand the correspondence between addresses for f and addresses for $\widetilde{f}$.

Given a point z in the f-plane, whose iterates always belong to $\mathcal{T}$, and such that $f^{n}(z) \in F_{\alpha_n, i_n}$, we define its *address* s as

$$s = \binom{\alpha_0}{i_0}\binom{\alpha_1}{i_1}\ldots .$$

The left-sided shift map σ is well defined, and a point with address s is mapped to a point with address σs.

Lemma 2.7. *A point* $\widetilde{z} \in J(\widetilde{f})$ *has address* $\widetilde{s} = \binom{\alpha_0}{i_0}\binom{\alpha_1}{i_1}\ldots$ *if and only if* $z = \exp(\widetilde{z})$ *has address* $s = \binom{\alpha_0}{i_1}\binom{\alpha_1}{i_2}\ldots$*.*

Proof. If $\widetilde{z} \in \widetilde{T}_{\alpha_n, i_n}$ and $\widetilde{f}(\widetilde{z}) \in \widetilde{T}_{\alpha_{n+1}, i_{n+1}} \subset \widetilde{S}_{i_{n+1}}$, then $\widetilde{z} \in \widetilde{F}_{\alpha_n, i_n, i_{n+1}}$ by the choice of lift $\widetilde{f}$. Hence $z = \exp(\widetilde{z}) \in F_{\alpha_n, i_{n+1}}$ by the choice of labeling for the fundamental domains. For the converse, observe that $\exp^{-1}(F_{\alpha_n, i_{n+1}}) = \{\widetilde{F}_{\alpha_n, m, i_{n+1}}\}_{m \in \mathbb{Z}}$ and $\widetilde{F}_{\alpha_n, m, i_{n+1}} \subset \widetilde{T}_{\alpha_n, m}$. □

Proposition 2.6 is a corollary of Theorem 2.4 and the following proposition:

Proposition 2.8 (**Lifting addresses**). *If* $s = \binom{\alpha_0}{i_0}\binom{\alpha_1}{i_1}...$ *is an address for* f*, there is a dynamic ray* g_s *which contains the image under the exponential map of any of the ray tails* $g_{\widetilde{s}_m}$*, where* $\widetilde{s}_m = \binom{\alpha_0}{m}\binom{\alpha_1}{i_0}...$*, and* $m \in \mathbb{Z}$*.*

Proof. The image of any of the dynamic ray tails $J_{\widetilde{s}_m}$ under the exponential map is a dynamic ray tail for f by definition, and it consists of points of address s by Lemma 2.7. □

We are only left to show that any ray is asymptotically contained in some fundamental domain.

Lemma 2.9. *Any dynamic ray is asymptotically contained in some fundamental domain.*

Proof. Suppose that a ray g is not asymptotically contained in any fundamental domain. Then for some sequence $t_n \to \infty$, $g(t_n)$ belongs to preimages of δ. Hence $f(g)$ intersects δ infinitely many times and $f^2(g)$ intersects D infinitely many times as $t \to \infty$, contradicting the definition of dynamic ray. □

This concludes the proof of part (b) in Proposition 2.2.

2.4 Expansion near infinity

We now prove that, when restricted to a finite number of fundamental domains and far enough to the right, $\widetilde{f}$ is uniformly expanding.

Proposition 2.10 (**Almost straight**). *Let* $\mathcal{F}$ *be a finite collection of fundamental domains. Then for any* R *sufficiently large, the preimage of the circle* C_R *of radius* R *restricted to* $\mathcal{F}$ *is contained in the bounded connected component of* $\mathbb{C} \setminus C_R$*.*

The remaining of this section is devoted to prove Proposition 2.10. We use the following result from Proposition 5.4 in [R3S]:

Proposition 2.11 (**Growth of real part**). *Let* $\widetilde{f}$ *be the lift of a function* f *in the class* $\widehat{\mathcal{B}}$*, and* $K > 1$*. Then there exist* $M > 0$ *such that the following holds: if* $\widetilde{T}$ *is a tract for* $\widetilde{f}$*, and* $u, v, \widetilde{f}(u), \widetilde{f}(v) \in \widetilde{T}$ *with* $\operatorname{Re}\widetilde{f}(u), \operatorname{Re}\widetilde{f}(v)$ *sufficiently large, and* $\operatorname{Re} v > \operatorname{Re} u + M$*, then* $\operatorname{Re}\widetilde{f}(v) \geq K \operatorname{Re}\widetilde{f}(u) + M$*.*

Again, the way in which these results are stated in [R3S] is somewhat different, so we include a sketch of the proof using their results and definitions.

Proof. By Corollary 5.8 in [R3S], lifts of functions in the class $\widehat{\mathcal{B}}$ satisfy statements (a) and (b) of Proposition 5.4 in [R3S]. Proposition 5.4 states that statements (a) and (b) are equivalent to statement (c) which is very similar to the statement of Proposition 2.11 here. There is an extra condition in statement (c) of Proposition 5.4 which is satisfied because by assuming $u, v, \widetilde{f}(u), \widetilde{f}(v) \in \widetilde{T}$, this condition becomes the definition of bounded slope (see Definition 5.1, always in [R3S]). Also, in [R3S], $\widetilde{f}$ is assumed to be normalized (see page 8 in [R3S]). This condition is replaced in our case by taking the real part of $f(u), f(v)$ to be sufficiently large. □

Before starting the proof of Proposition 2.10, let us recall that the density $\rho_{\mathbb{H}}(x)$ of the hyperbolic metric in the right half plane $\mathbb{H}$ is $\rho_{\mathbb{H}}(x) = \frac{1}{\operatorname{Re} x}$. Also, it is a standard estimate that the hyperbolic density on any simply connected hyperbolic domain Ω is comparable with the inverse of the distance to the boundary. More precisely,

$$\frac{1}{2\operatorname{dist}(x, \partial\Omega)} \leq \rho_\Omega(x) \leq \frac{2}{\operatorname{dist}(x, \partial\Omega)}.$$

As any tract $\widetilde{T}$ in logarithmic coordinates never has vertical width larger than 2π, the maximal distance of a point from the boundary is π, hence on the tract we have the inequality

$$\frac{1}{2\pi} \leq \rho_{\widetilde{T}}(x). \tag{2.1}$$

Therefore for any curve γ in $\widetilde{T}$, $\ell_{\mathrm{eucl}}(\gamma) \leq 2\pi\ell_{\widetilde{T}}(\gamma)$.

Proof of Proposition 2.10. Observe that it is enough to prove the claim for one fundamental domain F, and then for any finite family take R sufficiently large as to work for all of them. Recall the notation from Section 2.2, and let $\widetilde{f}$ be the lift of f defined on the tracts $\widetilde{F}$ with the conventions adopted there.

Let $F = F_{\alpha,i}$ be a fundamental domain for f, and let g_F be the fixed ray given by Proposition 2.6. Because g_F is a forward invariant curve asymptotically contained in F, its address is a fixed address s of the form $s = \overline{\binom{a}{i}}$. By Proposition 2.8, the lifts of g_F to the logarithmic coordinates is the family of rays with addresses $\widetilde{s}_m = \binom{a}{m}\overline{\binom{a}{i}}$. Let $\widetilde{F}$ be the fundamental domain for $\widetilde{f}$ which contains the ray tail $g_{\widetilde{F}}$ of address $\widetilde{s} = \overline{\binom{a}{i}}$. With this choice, $\widetilde{f}(g_{\widetilde{F}}) \subset g_{\widetilde{F}}$. Let us only consider points in $\widetilde{F}$ for which Proposition 2.11 holds. Observe that this is no real restriction, because it is $\widetilde{F}$ intersected with the preimage of a right half plane- which does not change anything on the geometry of $\widetilde{F}$ near infinity. Let $\widetilde{T}$ be the tract in which $\widetilde{F}$ is contained, and $\widetilde{S}_i$ be the strip which contains $\widetilde{T}$.

As $\widetilde{f} : \widetilde{T} \to \widetilde{H}$ is a biolomorphism, there is a well defined inverse $\widetilde{f}^{-1}_{\widetilde{T}}$, which is an isometry between the hyperbolic metric on $\widetilde{H}$ and the hyperbolic metric in $\widetilde{T}$. Its restriction $\widetilde{f}^{-1}_{\widetilde{F}} : \widetilde{S}_i \to \widetilde{F}$ is hence also an isometry between the hyperbolic metric on $\widetilde{H}$ and the hyperbolic metric in $\widetilde{T}$.

For any R, call L_R the vertical segment of length 2π and at real part R which joins two points on the curves $\widetilde{\delta}_i, \widetilde{\delta}_{i+1}$ bounding the strip $\widetilde{S}_i$ to which $\widetilde{T}$ belongs to. Call $\widetilde{S}_i^-$ the connected component of $\widetilde{S}_i \setminus L_R$ which is bounded (or, equivalently, which contains points with real part less than R). The image under the exponential map of L_R is a circle of radius e^R, which bounds the disk which contains the image of $\widetilde{S}_i^-$ under the exponential (see Figure 4).

So, using the fact that f is semi-conjugate to $\widetilde{f}$ on the tracts, it is enough to show that for sufficiently large R, $\widetilde{f}^{-1}_{\widetilde{F}}(L_R) \subset \widetilde{S}_i^-$.

Let $u \in g_{\widetilde{F}}$ such that $\operatorname{Re} \widetilde{f}(u) > \operatorname{Re} u$; many such points exist because for every $v \in g_{\widetilde{F}}$, $\operatorname{Re} f^n(v) \to \infty$. By Proposition 2.11, there is M such that for any $v \in g_{\widetilde{F}}$ with sufficiently large real part and with $\operatorname{Re} v > \operatorname{Re} u + M$, then $\operatorname{Re} \widetilde{f}(v) > \operatorname{Re} v + M$. For each such v,


$\widetilde{\delta}_{i+1}$
$\widetilde{T}_{\alpha,i}$
$f(v)$
v
$\widetilde{F}_{\alpha,i,i}$
$\widetilde{\delta}_i$
$L_r = L_{f(v)}$

Figure 4: Sketch of the proof of Proposition 2.10

consider the vertical segment $L_{\widetilde{f}(v)}$ of length 2π connecting the boundary of $\mathbb{S}_i$ and passing through $\widetilde{f}(v)$. Note that as v moves continuously along $g_{\widetilde{F}}$ with increasing real part, $\widetilde{f}(v)$ moves continuously as well satisfying $\operatorname{Re}\widetilde{f}(v) \geq \operatorname{Re} v + M$ (see Figure 4).

We will now show that as $\operatorname{Re} v \to \infty$, and in particular for $\operatorname{Re} v > R'$ with some $R' > 0$, the diameter of $\widetilde{f}_{\widetilde{F}}^{-1}(L_{\widetilde{f}(v)}) \to 0$, while $|v - \widetilde{f}(v)| \geq m$, so that $L_{\widetilde{f}(v)}$ and $\widetilde{f}_{\widetilde{F}}^{-1}(L_{\widetilde{f}(v)})$ do not intersect; this will prove the claim for $R = e^{R'}$.

Because $\rho_{\widetilde{H}}(x) \sim \frac{1}{\operatorname{Re} x}$, we have that

$$\operatorname{diam}_{\widetilde{H}}(L_{\widetilde{f}(v)}) \sim \frac{\pi}{\operatorname{Re}\widetilde{f}(v)},$$

so by holomorphicity of $\widetilde{f} : \widetilde{T} \to \widetilde{H}$,

$$\operatorname{diam}_{\widetilde{T}} \widetilde{f}_{\widetilde{F}}^{-1}(L_{\widetilde{f}(v)}) \sim \frac{\pi}{\operatorname{Re}\widetilde{f}(v)},$$

hence by Equation 2.1,

$$\operatorname{diam}_{\text{eucl}} \widetilde{f}_{\widetilde{F}}^{-1}(L_{\widetilde{f}(v)}) \leq 2\pi \operatorname{diam}_{\widetilde{T}} \widetilde{f}_{\widetilde{F}}^{-1}(L_{\widetilde{f}(v)}) \to 0 \text{ as } \operatorname{Re}\widetilde{f}(v) \to \infty.$$

As

$$|v - \widetilde{f}(v)| > \operatorname{Re}\widetilde{f}(v) - \operatorname{Re} v > M,$$

for $\operatorname{Re}\widetilde{f}(v)$ sufficiently large $L_{\widetilde{f}(v)} \cap \widetilde{f}_{\widetilde{F}}^{-1}(L_{\widetilde{f}(v)}) = \emptyset$. As $v \in \widetilde{f}_{\widetilde{F}}^{-1}(L_{\widetilde{f}(v)})$ and $v \in \widetilde{S}_i^-$, the claim follows. □

This concludes the proof of the Structural Lemma 2.2.

Remark 2.12. If L_R intersects $\partial\widetilde{S}_i$ in more than two points for $R = \operatorname{Re} f(v)$, all estimates still hold, because they are performed in the hyperbolic metric for $\widetilde{H}$ and $\widetilde{T}$ respectively. The only thing to do is to define the inverse branch of $\widetilde{f}$ so that $\widetilde{f}^{-1}(\widetilde{f}(v)) = v$ and extend it by continuity to L_R.

3 Location of interior fixed points

Let us now fix a function $f \in \widehat{\mathcal{B}}$, and restrict our discussion to fixed points and fixed rays. From now on, we assume that all fixed rays land.

We first prove a lemma about landing of fixed rays for fundamental domains which do not intersect D, and then collect some remarks about the structure of the plane in logarithmic coordinates and the distribution of the interior fixed points with respect to the partition of the plane induced by the fixed rays.

As tracts accumulate only at infinity, only finitely many fundamental domains can intersect D. On fundamental domains not intersecting the disk, the dynamics, and especially the fixed rays, are easy to study.

Proposition 3.1 (**Forced landing).** *If a fundamental domain F does not intersect D, it contains a unique fixed point w which is repelling, and the fixed ray g_F given by part* (*b*) *in the Structural Lemma 2.2 lands at w.*

Proof. Consider a circle C_R of radius R such that part (*c*) of the Structural Lemma 2.2 holds, and let D_R be the bounded connected component of $\mathbb{C} \setminus C_R$ (see Figure 5). As $f : F \to \mathbb{C} \setminus (\overline{D} \cup \delta)$ is univalent, there is a branch ψ of f^{-1} mapping $D_R \setminus (\overline{D} \cup \delta)$ to a proper subset of $F \cap D_R \setminus (\overline{D} \cup \delta)$ by the Structural Lemma part (c). By the Schwarz Lemma, ψ is strictly contracting and has a unique fixed point $w \subset F$. Hence, under iterates of ψ every point tends to a unique fixed point w which is repelling for f; in particular, any invariant curve converges to w. $\square$



Figure 5: If F does not intersect D, the fixed ray g_F lands at the unique fixed point in F which must be repelling. However fixed rays from other fundamental domains which do intersect D, can also land at the same fixed point. There are only a finite number of those freelanders though. Light shaded we see the image of the dark shaded domain.

Remark 3.2. Observe that the ray g_F does not necessarily land alone; other fixed rays coming from fundamental domains which do cross the disk might land together with g_F (see Figure 5).

Recall that a fixed point is an *interior fixed point* if there are no fixed rays landing at it.

Observe that interior fixed points cannot be in $\mathbb{C}\setminus(\overline{D}\cup\mathcal{T})$, because f maps $\mathbb{C}\setminus(\overline{D}\cup\mathcal{T})$ to D. Also, they cannot be in a fundamental domain not intersecting the disk by Proposition 3.1. So interior fixed points are contained either in D or in the finitely many fundamental domains which intersect D (see Figure 6).



Figure 6: Interior fixed points can only lie in the finite number of fundamental domains which intersect D, or in $D\setminus\overline{T}$.

Recall that Γ is the graph formed by fixed rays together with their endpoints, and that the connected regions of $\mathbb{C}\setminus\Gamma$ are called *basic regions* for f.

The following is a corollary of Proposition 3.1.

Corollary 3.3. *There are finitely many basic regions.*

Proof. As tracts do not accumulate in any compact set, there are only finitely many tracts intersecting D. As preimages of δ do not accumulate in a compact set either, there are finitely many fundamental domains intersecting D, hence by Proposition 3.1 all but finitely many rays must land in the interior of their respective fundamental domains. Therefore, only the finitely many remaining ones are free to land together with other rays. ☐

The next two lemmas are interesting but will not be used later on.

Lemma 3.4 (Crossing tracts). *A basic region cannot be fully contained inside a tract.*

Proof. This proof uses notation from Subsection 2.2. If a basic region is fully contained inside some tract T_α, there exists at least two fixed rays g^1 and g^2 landing at a common fixed point z which are completely contained in T_α. Let $\{z^k\}$ be the fiber of z under the exponential map, with the convention that $z^k\in\widetilde{S}_k$. Because z is fixed, there is a unique $z^*\in\{z^k\}$ which is fixed. Let $\widetilde{g}^1$ and $\widetilde{g}^2$ be the lifts of g^1, g^2 landing at z^*. It follows that these curves must be fixed ray tails for $\widetilde{f}$: indeed, the unique lifts of g^1, g^2 which are fixed are both forced to land at the unique fixed point. This gives two invariant ray tails for $\widetilde{f}$ which are contained in the same tract, giving a contradiction. ☐

Lemma 3.5. *Fixed ray pairs separate the set of singular values.*

Proof. If a ray pair does not separate the set of singular values, the logarithmic coordinates can be redefined so that D does not to intersect the ray pair. Then, the original ray pair is contained in $\mathbb{C} \setminus \overline{D}$, hence its preimages are fully contained inside tracts, contradicting Lemma 3.4. □

4 Index and Argument principle

To count the number of fixed points of f inside a basic region we will make use of the argument principle. For a more general theory about index see for example [Why].

For all of this section, let $\gamma : [a, b] \to \mathbb{C}$, with $a, b \in \mathbb{R}$, $a \leq b$, be a piecewise $\mathcal{C}^1$ curve, not necessarily closed unless specified. The symbol γ will denote both the function $\gamma(t)$ or the set $\gamma[a, b]$.

For a curve γ as above and a point $P \in \mathbb{C} \setminus \gamma$ we define the *index* of γ with respect to P as

$$\operatorname{Ind}(\gamma, P) := \operatorname{Re} \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - P} dz.$$

By definition, if γ is the union of consecutive arcs γ_i,

$$\operatorname{Ind}(\gamma, P) = \sum_i \operatorname{Ind}(\gamma_i, P).$$

Observe that if γ is closed, this is the usual definition of the winding number and therefore $\operatorname{Ind}(\gamma, P) \in \mathbb{Z}$. In that case, the index can be used to count the number of zeros of a map inside a Jordan curve.

Theorem 4.1 (Argument principle). *Let γ be a piecewise $\mathcal{C}^1$ Jordan curve bounding a region Ω. Let f be holomorphic in a neighborhood of $\overline{\Omega}$ such that $f(z) \neq 0$ for all points $z \in \gamma$. Let $Z(f)$ be the set of zeros of f. Then*

$$\operatorname{Ind}(f(\gamma), 0) = \#(Z(f) \cap \Omega)$$

counted with multiplicity.

For a parametrized curve $\gamma(t)$, we denote by $f(\gamma) - \gamma$ the curve defined by $f(\gamma(t)) - \gamma(t)$. The following is a corollary of the Argument Principle, and will be our main tool.

Corollary 4.2 (Counting fixed points). *Let γ be a piecewise $\mathcal{C}^1$ Jordan curve bounding a region Ω. Let f be holomorphic in a neighborhood of $\overline{\Omega}$ such that $f(z) \neq z$ for all points $z \in \gamma$. Let* Fixed(f) *be the set of fixed points of f. Then*

$$\operatorname{Ind}(f(\gamma) - \gamma, 0) = \#(\mathrm{Fixed(f)} \cap \Omega)$$

counted with multiplicity.

We will use the fact that calculating the index of a curve with respect to a point is invariant under homotopy in the following sense (see [Why], Theorem 3.1 in Chapter V).

Lemma 4.3 (Invariance under small perturbations). *If $\gamma, \widehat{\gamma}$ are piecewise $\mathcal{C}^1$ curves such that for any t, $|\gamma(t) - \widehat{\gamma}(t)| < |\gamma(t) - P|$ for some $P \in \mathbb{C}$, then*

$$\operatorname{Ind}(\gamma, P) = \operatorname{Ind}(\widehat{\gamma}, P).$$

In the proof of the main theorem, we will use Lemma 4.3 in the form of the three lemmas that we state and prove below (see Figure 7).

Lemma 4.4 (First Homotopy Lemma). *If $\gamma, \widehat{\gamma} : [0,1] \to \mathbb{C}$ are two piecewise $\mathcal{C}^1$ arcs such that $\gamma(0) = \widehat{\gamma}(0) = P_0$, $\gamma(1) = \widehat{\gamma}(1) = P_1$, and such that γ is homotopic to $\widehat{\gamma}$ relatively to a point P and keeping P_0, P_1 fixed, then*

$$\operatorname{Ind}(\gamma, P) = \operatorname{Ind}(\widehat{\gamma}, P).$$

Proof. Observe that there is a definite distance ϵ between P and the compact set formed by $\gamma, \widehat{\gamma}$ and all the curves in the homotopy between them. Recall also that every homotopy can be approximated by piecewise $\mathcal{C}^1$ homotopies. In particular, we can assume that the homotopy between γ and $\widehat{\gamma}$ always moves in steps smaller than ϵ among curves on which the index is well defined, hence locally constant by Lemma 4.3. $\square$

Lemma 4.5 (Second Homotopy Lemma). *Let γ be a piecewise $\mathcal{C}^1$ arc, f be a continuous function, and $f(\gamma)$ be a closed curve starting and ending at the same point P. Suppose $\gamma \cap f(\gamma) = \emptyset$. If $f(\gamma)$ is homotopic to P relatively to γ by a homotopy which keeps P fixed, then*

$$\operatorname{Ind}(f(\gamma) - \gamma, 0) = \operatorname{Ind}(P - \gamma, 0) = +\operatorname{Ind}(\gamma, P).$$

Proof. If $f(\gamma)$ is homotopic to P relatively to γ by a homotopy which keeps P fixed, then $f(\gamma) - \gamma$ is homotopic relatively to 0 to the curve $P - \gamma$, hence the first equality holds by Lemma 4.4 (observe that the homotopy keeps fixed the endpoints of the curve $f(\gamma) - \gamma$ because it keeps fixed both the endpoint of $f(\gamma)$ and the endpoints of γ). The second equality follows from the definition of index. Indeed,

$$\int_{P-\gamma} \frac{1}{z} dz = \int_a^b \frac{1}{P - \gamma(t)} (-\gamma'(t)) dt = \int_\gamma \frac{1}{z - P} dz.$$

$\square$

Lemma 4.6 (Third Homotopy Lemma). *Let γ is a simple piecewise $\mathcal{C}^1$ arc, and $f(\gamma)$ is a Jordan curve starting and ending at a point P such that*

- $\gamma \cap f(\gamma) = \emptyset$
- *γ is contained in the bounded connected component of $\mathbb{C} \setminus f(\gamma)$.*



Figure 7: From left to right, setup for the first, second and third homotopy lemas.

Then

$$\operatorname{Ind}(f(\gamma)-\gamma,0)=\operatorname{Ind}(\gamma,P)+N,$$

where N is the degree of $f|_\gamma : \gamma \to f(\gamma)$.

Proof. By Lemma 4.4, we can assume $f(\gamma)$ to a be a circle and γ to be a straight segment. The claim then follows by direct counting, for example using vectors joining $x \in \gamma$ to $f(x) \in f(\gamma)$. □

5 Global counting: proof of Theorem B

Recall that, given a fundamental domain F and its fixed dynamic ray g_F as given by the Structural Lemma 2.2, we say that g_F *lands alone* if it lands inside F and no other fixed ray lands there. By Proposition 3.1, there are only finitely many fundamental domains whose fixed rays do not land alone, which are contained in finitely many tracts T_α, say M tracts. For each such T_α, let $\mathcal{F}_\alpha$ be the collection of fundamental domains $F \subset T_\alpha$, whose fixed ray does not land alone or which are in between two fundamental domains whose fixed rays do not land alone.

Let $N_\alpha := \#\mathcal{F}_\alpha$, $N = \underset{\alpha}{\Sigma} N_\alpha$. We will refer as $\widehat{\mathcal{F}_\alpha}$ to the union of the fundamental domains contained in $\mathcal{F}_\alpha$.

From now on we will restrict the index α to these tracts, assuming that they are labeled consecutively respecting the cyclic order, and take the index α modulo M. Theorem B is a consequence of the following theorem.

Theorem 5.1 (Global counting). *Let A be the number of fundamental domains in $\bigcup_\alpha \mathcal{F}_\alpha$ whose fixed dynamic rays land alone. Then there are exactly $N-A+1$ fixed points which are either landing points of one of the remaining $N-A$ fixed rays, or interior fixed points.*

Because N is finite, up to redefining δ we can assume that δ does not intersect any fixed ray coming from $\underset{\alpha}{\cup}\widehat{\mathcal{F}_\alpha}$. Indeed, for each of these finitely many rays, the part which is not contained inside a tract is compact, hence we only have to avoid a finite number of non-intersecting Jordan arcs.

Let C_R be a circle of radius R given by the Structural Lemma 2.2 for the family of fundamental domains $\mathcal{F}_\alpha$, and r_α be the preimages of C_R restricted to $\mathcal{F}_\alpha$. As usual, give C_R and any other Jordan curve in this section a counterclockwise orientation; this induces an orientation on the arcs r_α (see Figure 8 for an illustration of this setup).

Let $\delta_\alpha^\pm$ be the arcs contained in the preimages of δ which bound the two fundamental domains in $\mathcal{F}_\alpha$ whose immediate neighbors do not belong to $\mathcal{F}_\alpha$, starting at the points $P_\alpha^\pm$ in which they intersect r_α and ending at the boundary of T_α. Label them so as to respect the cyclic order in the sense that δ_α^- comes before δ_α^+ and after $\delta_{\alpha-1}^+$.

Observe that $f(P_\alpha^\pm) = P$, where P belongs to the intersection of δ with C_R. Call δ_P the arc in δ connecting P to D.

Finally, let γ_α be Jordan arcs in $\mathbb{C} \setminus (\overline{\mathcal{T} \cup D})$ connecting $\delta_{\alpha-1}^+$ with δ_α^-, which do not intersect the set of fixed rays (this can be done, because only finitely many fixed rays intersect $\mathbb{C} \setminus \overline{\mathcal{T}}$ by Lemma 3.1).

Observe that one and exactly one of the γ_α intersects δ, because $\bigcup T_\alpha \cup D$ divides the plane into finitely many connected components, only one of which contains δ. Choose that specific γ_α, so as to intersect δ only once, on the sub-arc of δ connecting P to ∞. Note that the image of each γ_α is fully contained in D.

Call Γ_α the arcs $\delta_{\alpha-1}^+ \cup \gamma_\alpha \cup \delta_\alpha^-$ connecting $P_{\alpha-1}^+$ to P_α^-. We will count the number of fixed points for f inside the region $\widehat{U}$ enclosed by the Jordan curve

$$\Gamma = \bigcup_{\alpha \leq N} (\Gamma_\alpha \cup r_\alpha).$$

Observe that, for R large enough, all fixed points which are landing points of the fixed rays associated to the fundamental domains in $\mathcal{F}$ have to be contained in $\widehat{U}$. Also, because there are no fixed points in $\mathbb{C} \setminus (D \cup \mathcal{T})$, the final count does not depend on the choice of γ_α. We will also see that the result does not depend on C_R (and hence on the r_α) as long as C_R satisfies the Structural Lemma 2.2 (see Figure 8).

Proof of Theorem 5.1. Here we calculate $\mathrm{Ind}(f(\Gamma) - \Gamma, 0)$, by calculating it piece by piece on the r_α and on the Γ_α. Observe that by construction there are no fixed points on Γ. Let us first calculate $\mathrm{Ind}(f(r_\alpha) - r_\alpha, 0)$. Observe that $f(r_\alpha)$ is the circle C_R covered N_α times, starting and ending at P. Then by the Third homotopy Lemma 4.6,

$$\mathrm{Ind}(f(r_\alpha) - r_\alpha, 0) = N_\alpha + \mathrm{Ind}(r_\alpha, P).$$

Now let us calculate $\mathrm{Ind}(f(\Gamma_\alpha) - \Gamma_\alpha, 0)$. The image of each γ_α is a curve starting at P, moving along δ_P, staying inside $\overline{D}$ and then moving back to P along δ_P. Because D is a topological disk, it can be retracted to any point in its boundary, hence $f(\Gamma_\alpha)$ is homotopic to P by a homotopy keeping P fixed, inside the set $\overline{D} \cup \delta_P$ which does not intersect Γ_α (first retract $f(\gamma_\alpha)$ to the point $\delta_P \cap \overline{D}$, then slide it to P inside δ_P). Hence, by the Second homotopy Lemma 4.5,



Figure 8: Global counting. The curve Γ is the boundary of the shaded region $\widehat{U}$. Its image consists of the big circle C_R, together with δ_P and the images of the curves γ_i which are inside the disc. In this example the Argument Principle gives 8 fixed points inside the shaded region. Since there are only 7 rays, at least one of the fixed points must be an interior fixed point.

$$\operatorname{Ind}(f(\Gamma_\alpha) - \Gamma_\alpha, 0) = \operatorname{Ind}(\Gamma_\alpha, P).$$

Summing up, we obtain

$$\operatorname{Ind}(f(\Gamma) - \Gamma, 0) = \sum_\alpha \Big(\operatorname{Ind}(r_\alpha, P) + N_\alpha + \operatorname{Ind}(\Gamma_\alpha, P) \Big) = N + \operatorname{Ind}(\Gamma, P) = N + 1.$$

The last equality follows from the fact that Γ is a Jordan curve and P is contained in the connected component in the interior of Γ. Hence, there are exactly $N + 1$ fixed points in $\widehat{U}$ counted with multiplicity. □

Corollary 5.2. *If there is only one basic region, there is exactly one interior fixed point.*

Proof. The fact that there is only one basic region means that each of the N fixed rays coming from the fundamental domains in the $\mathcal{F}_\alpha$'s lands alone at a fixed point, accounting for N fixed points. Hence there is exactly one interior fixed point left from the counting. □

6 Local Counting. Proof of Theorem A

Let us now consider a basic region V and show that it contains exactly one interior fixed point. In the general case, the boundary of V contains fixed rays and fixed points, so the counting is slightly more articulated.

For the purpose of the proof in this section, the basic region V contains also all those rays landing alone. In other points, rays landing alone are NOT part of the boundary of V, which consists exclusively of ray pairs. Notice that we then need to prove that V contains as many fixed points as rays landing alone, plus one.

To apply the argument principle it is important not to have fixed points on the boundary of the domain. Hence, we first explain how to change the boundary of V near fixed points, in the two cases in which the fixed point in question is either repelling or parabolic; then, we define a smaller region $\widehat{V}$ on which to do the actual counting, and eventually perform the counting itself. Without loss of generality, up to redefining δ, we can assume that V does not intersect δ.

Fixed points on the boundary

Suppose $z_0 \in \partial V$ is a repelling or parabolic fixed point. We will slightly modify the boundary of V near z_0 making use of the local dynamics around z_0.

We start with the case of repelling fixed points. Consider z_0 to be a repelling fixed point on the boundary of V. There is a linearizing neighborhood L of z_0 on which the dynamics is conjugate to a linear expanding map; L is tiled by fundamental domains whose boundaries are Jordan curves.

Let g_1, g_2 be the two fixed rays landing at z_0 belonging to the boundary of V, so that we encounter first g_1 and then g_2 while moving along the boundary of V leaving V on the left.

Let γ be an arc on the boundary of a fundamental domain in $L \setminus V$ which joins a point $x_1 \in g_1$ to a point $x_2 \in g_2$; also let v_1 and v_2 be the vectors joining x_1 to $f(x_1)$, and x_2 to $f(x_2)$ respectively. Call Δ the increment counterclockwise (measured in turns, i.e. modulo 2π) going from v_1 to v_2 (modulo 2π); observe that going in the clockwise direction the increment is $-1 + (\Delta)$ (see Figure 9).

Lemma 6.1. *For a repelling fixed point, the index* $\mathrm{Ind}(f(\gamma) - \gamma, 0) = \Delta$*, where* γ *is the curve described above.*

Proof. Because the fundamental domain containing γ is mapped completely outside itself (to the next fundamental domain), the vector $f(\gamma(t)) - \gamma(t)$ just moves counterclockwise from v_1 to v_2. This can be computed directly using a schematic diagram (and, implicitly, the First and Second homotopy Lemmas 4.4 and 4.5). □

Observe that we could as well have made the region smaller so as to leave out the repelling fixed point. This would change the contribution by one full turn, which is the turn corresponding to the fixed point which has been left out.



Figure 9: The enlargement of the basic region when there is a repelling fixed point in the boundary.

We now consider the case of parabolic fixed points. The local dynamics are described by the Leau-Fatou flower theorem ([Mi], Theorem 10.7), which we state below. Petals, as defined in the introduction, are tiled by croissant-shaped *fundamental domains*, mapping to one another under the dynamics, whose boundaries are two Jordan arcs starting and ending at z_0.

Theorem 6.2 (Leau-Fatou Flower Theorem). *If z_0 is a parabolic point of multiplicity $m+1$, then there are exactly m repelling petals and m attracting petals attached z_0, alternating. Their union forms an open neighborhood of z_0.*

Now let us consider z_0 to be a parabolic fixed point on the boundary of V, with k attracting petals attached to z_0 and contained in V. Let g_1, g_2 be the two fixed rays landing at z_0 belonging to the boundary of V like before, x_1 and x_2 belonging to g_1, g_2 respectively and to a repelling petal for z_0. Observe that g_1, g_2 might land at z_0 from the same repelling petal, in which case $k = 0$ and x_1, x_2 belong to the same repelling petal. Again, let v_1 and v_2 be the vectors joining x_1 to $f(x_1)$, and x_2 to $f(x_2)$ respectively. See Figure 10.

Lemma 6.3. *We can join two points $x_1 \in g_1$ and $x_2 \in g_2$ inside V by a curve γ, such that* $\mathrm{Ind}(f(\gamma) - \gamma, 0) = -1 + \Delta - k$.

Proof. Let γ be a flower shaped curve constructed in the following way: start at x_1, move along the boundary of a fundamental domain inside the repelling petal until reaching an attracting petal; then move along the boundary of a fundamental domain inside the attracting petal until reaching the next repelling petal, and so on until reaching the ray g_2 at some point x_2. As before, let Δ be the increment counterclockwise between the vector v_1 joining x_1 to $f(x_1)$ and v_2 joining x_2 to $f(x_2)$. To calculate the index $\mathrm{Ind}(f(\gamma) - \gamma, 0)$ along γ, observe that the vector given by $f(\gamma) - \gamma$ turns clockwise until it reaches the attracting petal, then makes one full turn clockwise when moving inside the attracting petal (because now $f(\gamma)$ is inside γ), keeps moving clockwise along the repelling petal and so on, until it reaches the total increment of one full turn for each attracting petal inside, plus the total increment from



Figure 10: The reduction of the basic region when there is a parabolic point in the boundary. The sketch corresponds to one attracting petal in V ($k = 1$).

v_1 to v_2 clockwise. This gives an increment of $k + \Delta$ clockwise, which corresponds to an increment of $-1 + (\Delta) - k$ counterclockwise. □

When doing the counting, in addition to modifying the region near the fixed points, we will also *thicken* the fixed rays, i.e. we will consider a slightly larger region by substituting the fixed rays with two piecewise $\mathcal{C}^1$ curves near by whose image is outside the thickened region. This is a standard procedure and only relies on the fact that the dynamics is repelling near the fixed rays.

Proof of Theorem A

Let $\widehat{U}, \Gamma_\alpha, r_\alpha$ and $\mathcal{F}_\alpha$ be as in Section 5, and as before, consider the index α modulo M.

Proof of Theorem A. Fix a basic region V. Let us first suppose that there are no parabolic points on the boundary of V.

Let $\widehat{V}$ be the region obtained by intersecting V with $\widehat{U}$, thickening the rays and modified near the repelling fixed points as described in Lemma 6.1. Recall also that R was chosen large enough so that all fixed rays associated to the fundamental domains in $\mathcal{F}$ have their landing points in $\widehat{V}$. Let $\Gamma_{\widehat{V}}$ be the boundary of $\widehat{V}$. Up to redefining δ, we may assume that δ does not intersect $\widehat{V}$.

The only difference with respect to the global counting, is that we have to calculate the variation on the index when moving along one of the pieces of the thickened ray pairs forming $\Gamma_{\widehat{V}}$. Observe that there is one and exactly one thickened ray pair separating δ from $\widehat{V}$.

Let us keep the notation from Section 5, with one modification: let $\{P_i\}$ be the collection of all preimages of P intersecting $\Gamma_{\widehat{V}}$, labeled as to respect the cyclic order. Also, let s_i be the arc along $\Gamma_{\widehat{V}}$ joining two consecutive points P_i and P_{i+1} (see Figure 11).

The s_i can be of four types:

(0) s_i contains the thickened ray pair separating δ from $\widehat{V}$;

(1) s_i contains any other thickened ray pair;

(2) $s_i \subset r_\alpha$ for some r_α;

(3) $s_i = \Gamma_\alpha$ for some α.



Figure 11: An example of counting for the basic region shown in the picture. The shaded region is the subset which must contain the interior fixed points. Arcs of type (0) and (1) are colored in blue; arcs of type (2) are colored in green; arcs of type (3) are in red. Each arc has the same color as its image. In this example the index is 7 if we include the boundary fixed points.

Let N_1 be the number of arcs s_i of type (1), and N_2 be the number of arcs s_i of type (2).

We only have to compute $\mathrm{Ind}(f(s_i)-s_i, 0)$ when s_i is of type (0) and (1), as the other two cases are already covered in the previous section. In case (1), $f(s_i)$ is a Jordan curve starting at P, moving along C_R until it reaches the intersection with the thickened ray, moving along the ray pair until it reaches C_R again, and then moving back to P along C_R. Because the cyclic order is preserved, and using Lemma 4.6, we obtain that $\mathrm{Ind}(f(s_i) - s_i, 0) = 1 + \mathrm{Ind}(s_i, P)$.

In case (0), the same counting gives $\mathrm{Ind}(f(s_i) - s_i, 0) = 2 + \mathrm{Ind}(s_i, P)$. So in the three cases under consideration we obtain:

$$\begin{aligned} \mathrm{Ind}(f(s_i) - s_i, 0) &= 2 + \mathrm{Ind}(s_i, P) && \text{in case } (0); \\ \mathrm{Ind}(f(s_i) - s_i, 0) &= 1 + \mathrm{Ind}(s_i, P) && \text{in case } (1), (2); \\ \mathrm{Ind}(f(s_i) - s_i, 0) &= \mathrm{Ind}(s_i, P) && \text{in case } (3). \end{aligned}$$

Summing up, we have

$$\mathrm{Ind}(f(\Gamma_{\widehat{V}}) - \Gamma_{\widehat{V}}, 0) = N_1 + N_2 + 2 + \sum_{s_i} \mathrm{Ind}(s_i, P) = N_1 + N_2 + 2,$$

where we have used that P is not contained in the bounded connected component of $\mathbb{C} \setminus \Gamma_{\widehat{V}}$, so $\operatorname{Ind}(\Gamma_{\widehat{V}}, P) = 0$. We deduce from this that there is exactly one interior fixed point. Indeed,

- for the s_i of type (0), there is one fixed point in $\widehat{V}$ which was on the initial fixed ray pair.
- for each s_i of type (1), there is exactly one fixed point which was originally on the boundary of V and has been included in $\widehat{V}$ after the thickening procedure;
- for each s_i of type (2) there is exactly one fixed point in the interior $\widehat{V}$ which is the landing point of the unique fixed ray which is asymptotically contained in that fundamental domain (notice that those rays must land alone or otherwise they would be part of a ray pair).

This means, there are $N_1 + N_2 + 1$ fixed points in $\widehat{V}$ which are landing points of fixed rays; it follows that there is exactly one fixed point left, which has to be an interior fixed point because there are no more rays left to land on it.

Now replace n repelling periodic point on the boundary with n parabolic points, with k_i attracting petals contained in $\widehat{V}$. For each repelling periodic point replaced, by Lemmas 6.1 and 6.3 the counting changes by adding $-1 + \Delta - k_i$ and subtracting Δ. Putting everything together, we obtain that

$$\operatorname{Ind}(f(\Gamma_{\widehat{V}}) - \Gamma_{\widehat{V}}, 0) = N_1 + N_2 + 1 - n - \Sigma k_i.$$

Observe that there are at least $N_1 + N_2 + 1 - n$ fixed points inside the new region $\widehat{V}$ (the n parabolic points have now been left out of $\widehat{V}$), so it has to be $\operatorname{Ind}(f(\Gamma_{\widehat{V}}) - \Gamma_{\widehat{V}}, 0) \geq N_1 + N_2 + 1 - n$.

It follows that $\Sigma k_i \leq 1$, so that there is at most one attracting petal intersecting $\widehat{V}$ (however, there might be many parabolic points with no attracting petals in the initial region V). It follows that there are no interior fixed points if there is an attracting petal intersecting $\widehat{V}$, and that there is exactly one interior fixed point otherwise. □

7 Proof of corollaries D and E

As explained in the introduction, Theorem C is an immediate consequence of Theorem A, replacing f by f^p and observing that class $\widehat{\mathcal{B}}$ is closed under composition. In this section we prove some of the many corollaries of Theorem C. We recall that the period is never assumed to be minimal.

Corollary 7.1. *If f is a function in $\widehat{\mathcal{B}}$ whose periodic rays land, then*

1. *There are only finitely many interior periodic points of any given period.*
2. *Any two periodic Fatou components can be separated by a periodic ray pair.*
3. *There are no Cremer points on the boundary of periodic Fatou components.*

4. *If z_0 is a parabolic point, any two of its attracting petals are separated by a ray pair of the same period. In particular, for each repelling petal there is at least one ray which lands at z_0.*

5. *For any given period p, there are only finitely many (possibly none) periodic points of period p which are landing points of more than one periodic ray of period p. None of them is landing point of infinitely many rays of the same period.*

Proof of Theorem 7.1.

1. By Theorem C, for each p there are finitely many basic regions for f^p, and each of them contains at most an interior fixed point which is an interior periodic point of period p (non necessarily minimal) for f.

2. Let U_1, U_2 be two periodic Fatou components of least common period p. Then $\overline{U_1}$ and $\overline{U_2}$ contain each a point z_1, z_2 respectively which is fixed under f^p. If both z_1 and z_2 are in the interior of their Fatou component, they are interior periodic points, so they are separated by a pair of rays of period p by Theorem C. As dynamic rays cannot intersect Fatou components, and Fatou components are connected sets, it follows that the same ray pair separate U_1 from U_2. If instead at least one of them is on the boundary, say z_1, then z_1 is a parabolic point and U_1 contains an attracting petal for z_1. In particular, by Theorem C, there are no other interior periodic points of period p or attracting petals in the basic region V containing U_1, so U_2 has to be separated from U_1 by two rays on the boundary of V (observe that U_2 could also be another parabolic basin attached to the same parabolic point z_1).

3. Observe that Cremer points are interior periodic points, since by the Snail Lemma (see Lemma 16.2 in [Mi]) no periodic rays can land at Cremer points. Hence they can be separated from any other interior periodic point or periodic Fatou components as in the previous case.

4. If not, there would be two attracting petals belonging to the same basic region, contradicting Theorem C. The fact that each repelling petal contains at least one dynamic ray landing at z_0 then follows from the alternation between attracting and repelling petals given by the Flower Theorem 6.2.

5. If not, there would be infinitely many basic regions of period p, contradicting Theorem C.

□

From the fact that Fatou components can be separated by periodic ray pairs we obtain the following additional corollary. Given an invariant Siegel disk Δ, we say that U is a hidden component of Δ if U is a bounded connected component of $\mathbb{C} \setminus \overline{\Delta}$. Call $\widehat{\Delta} := \overline{\Delta} \cup (\bigcup U_i)$, where U_i are the hidden components for Δ. The proof of the next corollary follows the outline of Lemmas 2, 3 and 10 in [CR]; see also [Ro].

Corollary 7.2. *If $f \in \widehat{\mathcal{B}}$ has no wandering domains, any hidden component of a bounded invariant Siegel Disk is preperiodic to the Siegel Disk itself.*

Proof. Observe that $J(f)=\overline{I(f)}$: in fact, $I(f)$ and hence its closure are completely invariant, so $\overline{I(f)}\supset J(f)$, which is the smallest completely invariant set; on the other side, $\overline{I(f)}\subset J(f)$ by [EL]. Also, by Theorem 1.2 in [R3S], each $x\in I(f)$ can be joined to infinity by a curve of escaping points. Suppose that there is a hidden component U which is not preperiodic to Δ. Points in $\overline{\Delta}$ cannot be escaping because Δ is bounded and forward invariant. Since U is a bounded connected component of $\mathbb{C}\setminus\overline{\Delta}$, no point inside of U can be joined to infinity by a curve of escaping points. It follows that $U\cap J(f)=\emptyset$ and that U is a Fatou component. As $\partial U\subset J(f)$, it also follows that $\partial U\subset\partial\Delta$, hence $f(\partial U)\subset\partial\Delta$. Also, $\partial f(U)\subset f(\partial U)$ because f is holomorphic hence open, and $f(U)\cap\partial\Delta=\emptyset$ because $f(U)\subset F(f)$ and $\partial\Delta\subset J(f)$. Also $f(U)$ is bounded and connected because it is the image of a bounded connected set, and its boundary is contained in $\partial\Delta$, so $f(U)$ is also a hidden component of Δ. Hence, if there are no wandering domains, U is preperiodic to some periodic Fatou component $\widetilde{U}$ which is a hidden component of Δ; by part 4. of Corollary 7.1, unless $\widetilde{U}=\Delta$, they can be separated by two periodic rays landing together, hence $\partial\widetilde{U}\cap\partial\Delta$ is a single point- which is not possible as $\partial\widetilde{U}\subset\partial\Delta$. □

References

[Ba] Krzysztof Baranski, *Trees and hairs for some hyperbolic entire maps of finite order*, Math. Z. **257** (2007), 33–59.

[CR] Arnaud Chéritat, Pascale Roesch, *Herman's condition and Siegel disks of polynomials*, Preprint (2011), ArXiv:111.4629v1.

[DH] Adrien Douady, John H. Hubbard, *Etude dynamique des polynômes complexes*, Prépublications mathématiques d'Orsay (1984/1985), no. 2/4.

[DT] Robert L. Devaney, Folkert Tangerman, *Dynamics of entire functions near the essential singularity*, Ergodic Theory Dynam. Systems **6** (1986), no.4, 489–503.

[EL] Alexandre Eremenko, Mikhail Lyubich, *Dynamical properties of some classes of entire functions*, Ann. Inst. Fourier (Grenoble) **42** (1992), no. 4, 989–1020.

[Fa] Núria Fagella, *Dynamics of the complex standard family*, J. Math. Anal. Appl. **229** (1999), no.1, 1–31.

[GK] Lisa Goldberg, Linda Keen, *A finiteness theorem for a dynamical class of entire functions*, Ergodic Theory Dynam. Systems **6** (1986) no.2, 183–192.

[GM] Lisa Goldberg, John Milnor, *Fixed points of polynomial maps. Part II. Fixed point portraits*, Ann. Sci. École Norm. Sup. (4) **26** (1993), no. 1, 51–98.

[Mi] John Milnor, *Dynamics in one complex variable*, Annals of Mathematics Studies (2006), Princeton University Press.

[Rel] Lasse Rempe, *Siegel disks and periodic rays of entire functions*, J. Reine Angew. Math. **624** (2008), 81–102.

[R3S] Günter Rottenfusser, Johannes Rückert, Lasse Rempe, Dierk Schleicher, *Dynamic rays of bounded-type entire functions*, Annals of Mathematics **173** (2010),pp 77–125.

[Ro] James T. Rogers, *Diophantine conditions imply critical points on the boundaries of Siegel disks of polynomials*, Comm. Math. Phys. **195** (1), 175–193 (1998).

[Why] Gordon T. Whyburn, *Topological Analysis*, Princeton University Press (1958).